



Branchwidth of graphic matroids.

Frédéric Mazoit, Stéphan Thomassé

► To cite this version:

| Frédéric Mazoit, Stéphan Thomassé. Branchwidth of graphic matroids.. 2005. hal-00012312

HAL Id: hal-00012312

<https://hal.science/hal-00012312>

Preprint submitted on 19 Oct 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Branchwidth of graphic matroids.

Frédéric Mazoit* and Stéphan Thomassé†

Abstract

Answering a question of Geelen, Gerards, Robertson and Whittle [2], we prove that the branchwidth of a bridgeless graph is equal to the branchwidth of its cycle matroid. Our proof is based on branch-decompositions of hypergraphs.

1 Introduction.

Let $H = (V, E)$ be a hypergraph and (E_1, E_2) be a partition of E . The *border* of (E_1, E_2) is the set of vertices $\delta(E_1, E_2)$ which belong to both an edge of E_1 and an edge of E_2 . We often write it $\delta(E_1, E_2)$, or often simply $\delta(E_1)$. A *component* of E is a minimum nonempty subset $C \subseteq E$ such that $\delta(C) = \emptyset$. Let F be a subset of E . We denote by $c(F)$ the number of components of the subhypergraph of H spanned by F , i.e. the hypergraph $(V(F), F)$. A hypergraph H is *2-edge connected* if every vertex belongs to at least two edges and $c(E \setminus e) = 1$ for every $e \in E$.

A *branch-decomposition* \mathcal{T} of H is a ternary tree \mathcal{T} and a bijection from the set of leaves of \mathcal{T} into the set of edges of H . Practically, we simply identify the leaves of \mathcal{T} to the edges of H . Observe that every edge e of \mathcal{T} partitions $\mathcal{T} \setminus e$ into two subtrees, and thus correspond to a bipartition of E , called *e-separation*. More generally, a \mathcal{T} -separation is an *e-separation* for some edge e of \mathcal{T} . We will often identify the edge e of \mathcal{T} with the *e-separation*, allowing us to write, for instance, $\delta(e)$ instead of $\delta(E_1, E_2)$, where (E_1, E_2) is the *e-separation*.

Let f be a real function defined on the set of bipartitions of E . For sake of simplicity we often write $f(E_1)$ instead of $f(E_1, E \setminus E_1)$. Let \mathcal{T} be a branch-decomposition of H . The *f-width* of \mathcal{T} , denoted by $w_f(\mathcal{T})$, is the maximum value of $f(e)$, for all edges e of \mathcal{T} . The *f-branchwidth* of H , denoted by $\text{bw}_f(H)$, is the minimum *f-width* of a branch-decomposition of H . A branch-decomposition achieving $\text{bw}_f(H)$ is *f-optimal*.

The $|\delta|$ -branchwidth (i.e. when $f(E_1, E_2) = |\delta(E_1, E_2)|$) of a graph G is the usual branchwidth introduced by Robertson and Seymour in [5]. In this paper, we study the branchwidth associated to the function $\rho(E_1, E_2) = |\delta(E_1, E_2)| +$

*LIF - CMI, 39 rue Joliot-Curie, 13453 Marseille Cedex 13, France.
Frederic.Mazoit@lif.univ-mrs.fr

†Projet Mascotte, CNRS/INRIA/UNSA, INRIA Sophia-Antipolis, 2004 route des Lucioles
BP 93, 06902 Sophia-Antipolis Cedex, France. thomasse@univ-lyon1.fr

$2 - c(E_1) - c(E_2)$. Our goal is to prove that in the class of 2-edge connected hypergraphs, the $|\delta|$ -branchwidth is equal to the ρ -branchwidth. The proof simply consists to show that every 2-edge connected hypergraph admits a ρ -optimal decomposition such that $c(E_1) = c(E_2) = 1$ for every \mathcal{T} -separation (E_1, E_2) .

Our motivation comes from the following: Let M be a matroid on base set E with rank function r . The *weight* of every non-trivial partition (E_1, E_2) of E is $w(E_1, E_2) := r(E_1) + r(E_2) - r(E) + 1$. When \mathcal{T} is a branch-decomposition of M , i.e. a ternary tree whose leaves are labelled by E , the *width* of \mathcal{T} is the maximum weight of a \mathcal{T} -separation. Again, the *branchwidth* of M is the minimum width of a branch-decomposition of M . Let M be the cycle matroid of a 2-edge connected graph G , i.e. the matroid which base set is the set of edges of a graph and which independent sets are the acyclic subsets of edges. Branch-decompositions of G are exactly branch-decompositions of M . Moreover, $r(E_1) + r(E_2) - r(E) + 1$ is exactly $n_1 - c(E_1) + n_2 - c(E_2) - n + c(E) + 1$ where n_1, n_2, n are the number of vertices respectively spanned by E_1, E_2, E . Thus we have $w(E_1, E_2) = \delta(E_1) + 2 - c(E_1) - c(E_2) = \rho(E_1, E_2)$. In particular, the branchwidth of M is exactly the ρ -branchwidth, and thus is equal to the $|\delta|$ -branchwidth.

Unless stated otherwise, we always assume that H is a 2-edge connected hypergraph and \mathcal{T} is a branch-decomposition of H . Also, when speaking about width, branchwidth, etc, we implicitly mean ρ -width, ρ -branchwidth, etc.

2 Faithful branch-decompositions.

Let (E_1, E_2) be a \mathcal{T} -separation. The decomposition \mathcal{T} is *faithful* to E_1 if for every component C of E_1 , the partition $(C, E \setminus C)$ is a \mathcal{T} -separation. The *border graph* $G_{\mathcal{T}}$ has vertex set V and contains all edges xy for which there exists an edge e of \mathcal{T} such that $\{x, y\} \subseteq \delta(e)$. A branch-decomposition \mathcal{T}' is *tighter* than \mathcal{T} if $w_{\rho}(\mathcal{T}') < w_{\rho}(\mathcal{T})$ or if $w_{\rho}(\mathcal{T}) = w_{\rho}(\mathcal{T}')$ and $G_{\mathcal{T}'}$ is a subgraph of $G_{\mathcal{T}}$. Moreover, \mathcal{T}' is *strictly tighter* than \mathcal{T} if \mathcal{T}' is tighter than \mathcal{T} , and \mathcal{T} is not tighter than \mathcal{T}' . Finally, \mathcal{T} is *tight* if no \mathcal{T}' is strictly tighter than \mathcal{T} .

Lemma 1 *Let (E_1, E_2) be a partition of E . For any union E'_1 of connected components of E_1 and E_2 , we have both $\delta(E'_1) \subseteq \delta(E_1)$ and $\rho(E'_1) \leq \rho(E_1)$.*

□ Clearly, $\delta(E'_1) \subseteq \delta(E_1)$. Moreover, every vertex of $\delta(E_1)$ belongs to one component of E_1 and one component of E_2 . Therefore, if C is a component of E'_1 which is the union of k components of E_1 and E_2 , there are at least $k - 1$ vertices of $C \setminus \delta(C)$ which belong to $\delta(E_1)$. In all, the weight of the separation increased by $k - 1$ since we merge k components into one, but it also decreased by at least $k - 1$ since we lose at least that many vertices on the border. Since this is the case for every component of E'_1 or of $E \setminus E'_1$, we have $\rho(E'_1) \leq \rho(E_1)$. ■

Lemma 2 *Let (E_1, E_2) be an e -separation of \mathcal{T} . Let \mathcal{T}_1 be the subtree of $\mathcal{T} \setminus e$ with set of leaves E_1 . If \mathcal{T} is not faithful to E_1 , one can modify \mathcal{T}_1 in \mathcal{T} to form a tighter branch-decomposition \mathcal{T}' of H .*

□ Fix the vertex $e \cap \mathcal{T}_1$ as a root of \mathcal{T}_1 . Our goal is to change the binary rooted tree \mathcal{T}_1 into another binary rooted tree \mathcal{T}'_1 . For every connected component C of E_1 , consider the subtree \mathcal{T}_C of \mathcal{T}_1 which contains the root of \mathcal{T}_1 and has set of leaves C . Observe that \mathcal{T}_C is not necessarily binary since \mathcal{T}_C may contain paths having internal vertices with only one descendant. We simply replace these paths by edges to obtain our rooted tree \mathcal{T}'_C . Now, consider any rooted binary tree BT with $c(E_1)$ leaves and identify these leaves to the roots of \mathcal{T}'_C , for all components C of E_1 . This rooted binary tree is our \mathcal{T}'_1 . We denote by \mathcal{T}' the branch-decomposition we obtain from \mathcal{T} by replacing \mathcal{T}_1 by \mathcal{T}'_1 . Roughly speaking, we merged all subtrees of \mathcal{T}_1 induced by the components of E_1 together with $\mathcal{T} \setminus \mathcal{T}_1$ to form \mathcal{T}' . Let us prove that \mathcal{T}' is tighter than \mathcal{T} . For this, consider an edge f' of \mathcal{T}' . If $f' \notin \mathcal{T}'_1$, the f' -separations of \mathcal{T} and \mathcal{T}' are the same. If $f' \in BT$, by Lemma 1, we have $\rho(f') \leq \rho(e)$ and $\delta(f') \subseteq \delta(e)$. So the only case we have to care of is when f' is an edge of some tree \mathcal{T}'_C , where C is a component of E_1 . Recall that f' corresponds to a path P of \mathcal{T}_C . Let f be any edge of P . Let $(F, E \setminus F)$ be the f -separation of \mathcal{T} , where $F \subseteq E_1$. Therefore, the f' -separation of \mathcal{T}' is $(F \cap C, E \setminus (F \cap C))$. Since F is a subset of E_1 , the connected components of F are subsets of the connected components of E_1 . Thus $F \cap C$ is a union of connected components of F . This implies that $\delta(f') \subseteq \delta(f)$. Also, by Lemma 1, $\rho(f') \leq \rho(f)$.

We have proved that $w(\mathcal{T}') \leq w(\mathcal{T})$ and that $G_{\mathcal{T}'}$ is a subgraph of $G_{\mathcal{T}}$, thus \mathcal{T}' is tighter than \mathcal{T} . ■

3 Connected branch-decompositions.

Let $F \subseteq E$ be a component. The hypergraph $H * F$ on vertex set V and edge set $(E \setminus F) \cup \{V(F)\}$ is denoted by $H * F$. In other words, $H * F$ is obtained by merging the edges of F into one edge. A partition (E_1, E_2) of E is *connected* if $c(E_1) = c(E_2) = 1$. A branch-decomposition \mathcal{T} is *connected* if all its \mathcal{T} -separations are connected.

Lemma 3 *If \mathcal{T} is tight, every \mathcal{T} -separation (E_1, E_2) is such that E_1 or E_2 is connected.*

□ Suppose for contradiction that there exists a \mathcal{T} -separation (E_1, E_2) such that neither E_1 nor E_2 is connected. By Lemma 2, we can assume that \mathcal{T} is faithful to E_1 and to E_2 . Let \mathcal{C}_1 and \mathcal{C}_2 be respectively the sets of components of E_1 and E_2 . Consider the graph on set of vertices $\mathcal{C}_1 \cup \mathcal{C}_2$ where $C_1 C_2$ is an edge whenever $C_1 \in \mathcal{C}_1$ and $C_2 \in \mathcal{C}_2$ have nonempty intersection. This graph is connected and is not a star. Thus, it has a vertex-partition into two connected subgraphs, each having at least two vertices. This vertex-partition corresponds to a partition $(\mathcal{C}'_1, \mathcal{C}'_2)$ of $\mathcal{C}_1 \cup \mathcal{C}_2$.

Consider any rooted binary tree BT with $|\mathcal{C}'_1|$ leaves. Since every $C \in \mathcal{C}'_1$ is an element of $\mathcal{C}_1 \cup \mathcal{C}_2$, $(C, E \setminus C)$ is an e -separation of \mathcal{T} . We denote by \mathcal{T}_C the tree of $\mathcal{T} \setminus e$ with set of leaves C . Root \mathcal{T}_C with the vertex $e \cap \mathcal{T}_C$ in order to get a binary rooted tree. Now identify the leaves of BT with the roots of \mathcal{T}_C , for $C \in \mathcal{C}'_1$. This rooted tree is our \mathcal{T}'_1 . We construct similarly \mathcal{T}'_2 . Adding an edge between the roots of \mathcal{T}'_1 and \mathcal{T}'_2 gives the branch-decomposition \mathcal{T}' of H . By Lemma 1, $w(\mathcal{T}') \leq w(\mathcal{T})$. Moreover, $G_{\mathcal{T}'}$ is a subgraph of $G_{\mathcal{T}}$. Let us now show that $G_{\mathcal{T}'}$ is a strict subgraph of $G_{\mathcal{T}}$. Indeed, since \mathcal{C}'_1 is connected and has at least two elements, it contains $C_1 \in \mathcal{C}_1$ and $C_2 \in \mathcal{C}_2$ such that $C_1 \cap C_2$ is nonempty. By construction, every vertex x of $C_1 \cap C_2$ is such that $x \notin \delta(\mathcal{C}'_1)$ and $x \in \delta(\mathcal{C}_1)$. Similarly, there is a vertex y spanned by \mathcal{C}'_2 such that $y \notin \delta(\mathcal{C}'_2)$ and $y \in \delta(\mathcal{C}_2)$. Thus xy is an edge of $G_{\mathcal{T}}$ but not of $G_{\mathcal{T}'}$, contradicting the fact that \mathcal{T} is tight. \blacksquare

Theorem 1 *For every branch-decomposition \mathcal{T} of a hypergraph H , there exists a tighter branch-decomposition \mathcal{T}' such that for every \mathcal{T}' -separation (E_1, E_2) with $c(E_1) > 1$, E_1 consists of components of $H \setminus e$, for some $e \in E$. In particular, if H is 2-edge connected, it has an optimal connected branch-decomposition.*

\square Let us prove the theorem by induction on $|V| + |E|$. The statement is obvious if $|E| \leq 3$, so we assume now that H has at least four edges. Call *achieved* a branch-decomposition satisfying the conclusion of Theorem 1. If \mathcal{T} is not tight, we can replace it by a tight branch-decomposition tighter than \mathcal{T} . So we may assume that \mathcal{T} is tight.

If H is not connected, apply induction on every components of H in order to find an achieved branch-decomposition. Then merge these branch-decompositions into one branch-decomposition of H .

If there is an edge $e \in E$ such that $H \setminus e$ is not connected, we can assume by Lemma 2 that \mathcal{T} is faithful to $E \setminus e$. Let E_1 be a connected component of $E \setminus e$. Let \mathcal{T}_1 be the branch-decomposition induced by \mathcal{T} on $E_1 \cup e$. Let also \mathcal{T}_2 be the branch-decomposition induced by \mathcal{T} on $(E \setminus E_1) \cup e$. By the induction hypothesis, there exists two achieved branch-decompositions \mathcal{T}'_1 and \mathcal{T}'_2 , respectively tighter than \mathcal{T}_1 and \mathcal{T}_2 . Identify the leaf e of the trees \mathcal{T}'_1 and \mathcal{T}'_2 , and attach a leaf labelled by e to the identified vertex. Call \mathcal{T}' this branch-decomposition of H . Observe that it is tighter than \mathcal{T} and achieved.

So we assume now that H is 2-edge connected. The key-observation is that if there is a connected \mathcal{T} -separation (E_1, E_2) with $|E_1| \geq 2$ and $|E_2| \geq 2$, we can apply the induction hypothesis on $H * E_1$ and $H * E_2$ and merge the two branch-decompositions to obtain an optimal connected branch-decomposition of H . Therefore, we assume that every \mathcal{T} -separation (E_1, E_2) with $|E_1| \geq 2$ and $|E_2| \geq 2$ is such that E_1 or E_2 is connected.

We now orient the edges of \mathcal{T} . If (E_1, E_2) is an e -separation such that E_2 is connected but not an edge of H , we orient e from E_1 to E_2 . Since H is 2-edge-connected, every edge of \mathcal{T} incident to a leaf is oriented from the leaf. By Lemma 3, every edge get at least one orientation. And by the key-observation, every edge of \mathcal{T} has exactly one orientation.

This orientation of \mathcal{T} has no circuit, thus there is a vertex $t \in T$ with outdegree zero. Since every leaf has outdegree one, t has indegree three. Let us denote by A, B, C the set of leaves of the three trees of $\mathcal{T} \setminus t$. Observe that by construction, $A \cup B$, $A \cup C$ and $B \cup C$ are connected. By Lemma 2, we can assume moreover that \mathcal{T} is faithful to A, B and C . We claim that A is a disjoint union of edges, i.e. the connected components of A are edges of H . To see this, pick any component C_A of A . Since \mathcal{T} is faithful to A , $(C_A, E \setminus C_A)$ is a \mathcal{T} -separation. But this is simply impossible since $B \cup C$ being connected, $E \setminus C_A$ is also connected, against the fact that every edge has a single orientation. So the hypergraph H consists of three sets of disjoint edges A, B, C . Call this partition the *canonical partition* of \mathcal{T} . Call $(A, E \setminus A)$, $(B, E \setminus B)$ and $(C, E \setminus C)$ the *main* \mathcal{T} -separations. The width of every other \mathcal{T} -separation is strictly less than $\text{bw}(H)$. Since every vertex of H belongs to two or three edges, it is spanned by at least two of the sets $\delta(A), \delta(B), \delta(C)$. In particular $G_{\mathcal{T}}$ is a complete graph, and thus every optimal branch-decomposition of H is tighter than \mathcal{T} . Set $\delta_{AB} := |\delta(A) \cap \delta(B)|$, $\delta_{AC} := |\delta(A) \cap \delta(C)|$, $\delta_{BC} := |\delta(B) \cap \delta(C)|$ and $\delta_{ABC} := |\delta(A) \cap \delta(B) \cap \delta(C)|$. We now prove some properties of H .

1. Two of the sets A, B, C have at least two edges. Indeed, assume for contradiction that $A = \{a\}$ and $B = \{b\}$. Since $|E| \geq 4$, there are at least two edges in C . Let $c \in C$. Assume without loss of generality that $|a \cap c| \geq |b \cap c|$. Now form a new branch-decomposition \mathcal{T}' by *moving* c to A , i.e. \mathcal{T}' has a separation $(A \cup c, B \cup (C \setminus c))$ and then four branches $A, c, B, (C \setminus c)$. We have

$$\rho(A \cup c, B \cup (C \setminus c)) = \rho(A) + |b \cap c| - |a \cap c|$$

since both parts are connected. In particular \mathcal{T}' is tighter than \mathcal{T} , and since the \mathcal{T}' -separation $(A \cup c, B \cup (C \setminus c))$ is connected and both of its branches have at least two vertices, we can apply induction to conclude.

2. Every set A, B, C have at least two edges. Indeed, assume for contradiction that A consists of a single edge a . Let b be an edge of B . If $|b \cap \delta(C)| \leq |b \cap a|$, we can as previously move b to A in order to conclude. Call $|b \cap \delta(C)| - |b \cap a|$ the *excess* of b . Similarly, call $|c \cap \delta(B)| - |c \cap a|$ the *excess* of an edge $c \in C$. Let s be the minimum excess of an edge e_s of $B \cup C$. Observe that $s \geq 1$ and that every $b \in B$ satisfies $|b \cap \delta(C)| \geq |b \cap \delta(A)| + s$. Thus, summing for all edges of B , we obtain $\delta_{BC} \geq \delta_{AB} + s|B|$. Similarly, $\delta_{BC} \geq \delta_{AC} + s|C|$. Note also that $\text{bw}(H) \geq \rho(C) = \delta_{BC} + \delta_{AC} - \delta_{ABC} - |C| + 1$ and $\text{bw}(H) \geq \rho(B) = \delta_{BC} + \delta_{AB} - \delta_{ABC} - |B| + 1$. In all

$$2 \text{bw}(H) \geq 2\delta_{BC} - 2\delta_{ABC} + \delta_{AC} - |C| + \delta_{AB} - |B| + 2.$$

Then $2 \text{bw}(H) \geq \delta_{AB} + s|B| + \delta_{AC} + s|C| - 2\delta_{ABC} + \delta_{AC} - |C| + \delta_{AB} - |B| + 2$. Finally, $\text{bw}(H) \geq \delta_{AC} + \delta_{AB} - \delta_{ABC} + 1 + ((s-1)|C| + (s-1)|B|)/2$. Since $\rho(A) = \delta_{AC} + \delta_{AB} - \delta_{ABC}$, we have $\text{bw}(H) \geq \rho(A) + s$. But then we can move e_s to A to conclude.

3. We have $\text{bw}(H) = \rho(A)$. If not, pick two edges a, a' of A and merge them together. The hypergraph we obtain is still 2-edge-connected, and the branch-decomposition still has the same width. Apply induction to get an achieved branch-decomposition. Then replace the merged edge by the two original edges. This branch-decomposition \mathcal{T}' is optimal, hence tighter than \mathcal{T} . So either we can apply induction on \mathcal{T}' , or \mathcal{T}' has a canonical partition. But in this last case, the canonical partition of \mathcal{T}' is exactly $\{a\}, \{a'\}, E \setminus \{a, a'\}$. Thus by 1, we can apply induction. Similarly, $\text{bw}(H) = \rho(B) = \rho(C)$.
4. We have $\text{bw}(H) \geq \beta + 1$, where β is the size of a maximum edge e of H . Since edges of H with only one vertex play no role here, we can ignore them. So the size of an edge of H is at least two. Assume for instance that $e \in A$. Since A has at least two components, we have $\text{bw}(H) = \rho(A) \geq |\delta(A)| - c(B \cup C) - c(A) + 2 \geq 2 + \beta - 1$.
5. We have $\delta_{ABC} = 0$. Indeed, suppose for contradiction that there exists a vertex z in $\delta(A) \cap \delta(B) \cap \delta(C)$. Consider the hypergraph H_z obtained from H by removing the vertex z from all its edges. The branch-decomposition \mathcal{T} induces a branch-decomposition \mathcal{T}_z of H_z having width at most $\text{bw}(\mathcal{T}) - 1$. Observe also that H_z is connected since z is incident to three edges and H is 2-edge connected. We apply induction on \mathcal{T}_z to obtain an achieved branch-decomposition \mathcal{T}'_z of H_z . Now add back the vertex z to the edges of H_z and call \mathcal{T}' the branch-decomposition obtained from \mathcal{T}'_z . Observe that if a \mathcal{T}'_z -separation (E_1, E_2) is connected, adding z will raise by at most one its width in \mathcal{T}' . Moreover if a \mathcal{T}'_z -separation (E_1, E_2) is not connected, say $c(E_2) > 1$, adding z can raise by at most two its width in \mathcal{T}' (either by merging three components of E_2 into one, or by merging two and increasing the border by one). Since \mathcal{T}'_z is achieved, E_2 is a set of components of $E \setminus e$ for some edge e of E . But then, $\rho_{\mathcal{T}'_z}(E_1, E_2) \leq |e| - 3 + 2 \leq \beta - 1 \leq \text{bw}(H) - 2$, and thus $\rho_{\mathcal{T}'}(E_1, E_2) \leq \text{bw}(H)$. Finally \mathcal{T}' is optimal. Moreover every \mathcal{T}' -separation (E_1, E_2) is connected. Indeed, if E_1 is connected in \mathcal{T}'_z , we are done. If E_1 is not connected in \mathcal{T}'_z , E_1 consists of components of $H_z \setminus e$, for some edge e of H_z . But since H is 2-edge connected, every component of E_1 in H contains z , otherwise they would be components of $H \setminus e$. Consequently E_1 is connected.
6. Every edge of H is incident to at least four other edges. Indeed, assume for contradiction that an edge a of A is incident to only one edge b of B and at most two edges of C . Moving a to B increases $\rho(B)$ by $|a \cap \delta(C)| - |a \cap b|$ and does not increase $\rho(A)$ and $\rho(C)$. Therefore, if $|a \cap \delta(C)| \leq |a \cap b|$, we can move a to B , and this new branch-decomposition \mathcal{T}' is strictly tighter than \mathcal{T} since the vertices of $a \cap b$ are no more joined to $(\delta(A) \setminus a) \cap \delta(C)$ in the graph $G_{\mathcal{T}'}$. Thus $|a \cap \delta(C)| \geq |a \cap b| + 1$. Moreover, moving a to C , increases $\rho(C)$ by at most $|a \cap b| - |a \cap \delta(C)| + 1$, since at most two components of C can merge. So $|a \cap b| + 1 > |a \cap \delta(C)|$, a contradiction.

This implies in particular that $\rho(e) \leq \text{bw}(H) - 3$ whenever e is not one of the main \mathcal{T} -separations. In particular $\beta \leq \text{bw}(H) - 3$.

7. The hypergraph H is triangle-free. Indeed, suppose that there exists three edges $a \in A$, $b \in B$ and $c \in C$ and three vertices $x \in a \cap b$, $y \in b \cap c$ and $z \in c \cap a$. Let H_{xyz} be the hypergraph obtained by removing x, y, z in the vertex set of H and in every edge of H . The branchwidth of H_{xyz} is at most $\text{bw}(H) - 2$, since we removed two vertices in the border of every main separation. As in 5, \mathcal{T} induces a branch-decomposition \mathcal{T}_{xyz} of H_{xyz} which can be improved by induction to an achieved branch-decomposition \mathcal{T}'_{xyz} of H'_{xyz} . Adding back the vertices x, y, z , we obtain a branch-decomposition \mathcal{T}' of H . We claim that $w(\mathcal{T}') \leq \text{bw}(H)$. Let (E_1, E_2) be a \mathcal{T}' -separation, without loss of generality, we assume that E_1 contains at least two edges of a, b, c , say a and b . Assume first that (E_1, E_2) is connected in \mathcal{T}'_{xyz} . If $c \in E_1$, we have $\rho_{\mathcal{T}'_{xyz}}(E_1) = \rho_{\mathcal{T}'}(E_1)$. If $c \notin E_1$, $\rho_{\mathcal{T}'_{xyz}}(E_1) + 2 = \rho_{\mathcal{T}'}(E_1)$ since we add the endvertices of c to the border of E_1 . Now, if (E_1, E_2) is not connected in \mathcal{T}'_{xyz} , then $\rho_{\mathcal{T}'_{xyz}}(E_1)$ is at most the size of an edge of H_{xyz} , hence at most β . Since x, y, z have degree two in H , each can either increase the border of a separation by one, or merge two components. In all, $\rho_{\mathcal{T}'_{xyz}}(E_1)$ increases by at most three. Since $\beta \leq \text{bw}(H) - 3$, we have that $w(\mathcal{T}') \leq \text{bw}(H)$. To conclude, we prove that \mathcal{T}' is connected. Indeed, if $c(E_2) > 1$ for some \mathcal{T}' -separation (E_1, E_2) , E_2 consists of components of $H_{xyz} \setminus e$ for some edge e of H_{xyz} . Since H is 2-edge connected, these components are not components of $H \setminus e$, so each of them must contain one of the edges a, b or c , and therefore E_2 is connected in \mathcal{T}' .

Now we are ready to finish the proof. Note that $\text{bw}(H) = (\rho(A) + \rho(B) + \rho(C))/3 = (2|V| - |E|)/3 + 1$. Consider the line multigraph $L(H)$ of H , i.e. the multigraph on vertex set $A \cup B \cup C$ and edge set V such that $v \in V$ is the edge which joins the two edges e, f of H such that $v \in e$ and $v \in f$. The multigraph $L(H)$ satisfies the hypothesis of Lemma 4 (proved in the next section), thus it admits a vertex-partition of its vertices as in the conclusion of Lemma 4. This corresponds to a partition of $A \cup B \cup C$ into two subsets $E_1 := A_1 \cup B_1 \cup C_1$ and $E_2 := A_2 \cup B_2 \cup C_2$ such that $|\delta(E_1, E_2)| \leq (2|V| - |E|/3) + 1$ and both E_1 and E_2 have at least $\lfloor |E|/2 \rfloor - 1$ internal vertices. In particular, the separation (E_1, E_2) has width at most $\text{bw}(H)$. Let us show that one of $\rho(A_1 \cup B_1)$, $\rho(B_1 \cup C_1)$, and $\rho(C_1 \cup A_1)$ is also at most $\text{bw}(H)$. For this, observe that

$$\delta(A_1 \cup B_1) + \delta(B_1 \cup C_1) + \delta(C_1 \cup A_1) \leq 2(|V| - (|E_2| - |\delta(E_2)|)).$$

Thus $\delta(A_1 \cup B_1) + \delta(B_1 \cup C_1) + \delta(C_1 \cup A_1) \leq 2|V| - 2\lfloor |E|/2 \rfloor + 2 \leq 2|V| - |E| + 3$. Without loss of generality, we can assume that $\delta(A_1 \cup B_1) \leq (2|V| - |E|)/3 + 1 = \text{bw}(H)$, and thus we split E_1 into two branches $A_1 \cup B_1$ and C_1 . We similarly split E_2 to obtain an optimal branch-decomposition \mathcal{T}' of H . Observe that since both $E_1 \setminus \delta(E_1)$ and $E_2 \setminus \delta(E_2)$ are not empty, the graph $G_{\mathcal{T}'}$ is not complete, against the fact that \mathcal{T} is tight, a contradiction. \blacksquare

4 The technical Lemma.

Let G be a multigraph and X, Y two subsets of its vertices. We denote by $e(X, Y)$ the number of edges of G between X and Y . We also denote by $e(X)$ the number of edges in X .

Lemma 4 *Let G be a 2-connected triangle-free multigraph on $n \geq 5$ vertices and m edges. Assume that its minimum underlying degree (forgetting the multiplicity of edges) is four. Assume moreover that its maximum degree is at most $(2m - n)/3 + 1$. There exists a partition (X, Y) of the vertex set of G such that $e(X) \geq \lfloor n/2 \rfloor - 1$, $e(Y) \geq \lfloor n/2 \rfloor - 1$ and $e(X, Y) \leq (2m - n)/3 + 1$.*

□ Call *good* a partition which satisfies the conclusion of Lemma 4. Assume first that there are vertices x, y such that $e(x, y) \geq \lfloor n/2 \rfloor - 1$. The minimum degree in $V \setminus \{x, y\}$ is at least two, so $e(V \setminus \{x, y\})$ is at least $\lfloor n/2 \rfloor - 1$. Thus, if the partition $(V \setminus \{x, y\}, \{x, y\})$ is not good, we necessarily have $d(x) + d(y) - 2e(x, y) > (2m - n)/3 + 1$. By the maximum degree hypothesis, both $d(x)$ and $d(y)$ are greater than $2e(x, y)$. Since G is triangle-free, there exists a partition (X, Y) where $(N(x) \cup x) \setminus y \subseteq X$ and $(N(y) \cup y) \setminus x \subseteq Y$. Observe that $e(X) \geq d(x) - e(x, y) > e(x, y) \geq \lfloor n/2 \rfloor - 1$. Similarly $e(Y) \geq \lfloor n/2 \rfloor - 1$. Moreover, since $m \geq 2n$ by the minimum degree four hypothesis, we have

$$\begin{aligned} e(X, Y) &\leq m - (d(x) + d(y) - 2e(x, y)) \\ &< m - (2m - n)/3 - 1 \\ &\leq (m + n)/3 - 1 \\ &\leq (2m - n)/3 + 1. \end{aligned}$$

So (X, Y) is a good partition. We assume from now on that the multiplicity of an edge is less than $\lfloor n/2 \rfloor - 1$.

Let $a + b = n$, where $a \leq b$. A partition (X, Y) of V is an *a-partition* if $|X| \leq a$, $e(X) \geq a - 1$, $e(Y) \geq b - 1$, $e(X, Y) \leq (2m - n)/3 + 1$, and the additional requirement that X contains a vertex of G with maximum degree.

Note that there exists a 1-partition, just consider for this $X := \{x\}$, where x has maximum degree in G (the minimum degree in Y is at least three, insuring that $e(Y) \geq n - 2$). We consider now an *a-partition* (X, Y) with maximum a . If $a \geq b - 1$, this partition is good and we are done. So we assume that $a < b - 1$. In particular $e(X) = a - 1$. The *excess* of a vertex $y \in Y$ is $exc(y) := d_Y(y) - d_X(y)$.

The key-observation is that there exists at most one vertex $y \in Y$ such that $e(Y \setminus y) < b - 2$. Indeed, if there is a vertex of Y with degree one in Y , we simply move it to X , and we obtain an $(a + 1)$ -partition ($e(X)$ increases, $e(Y)$ decreases by one, and $e(X, Y)$ decreases). Thus the minimum degree in Y is at least two. Moreover, if there is a vertex of Y with degree two, we can still move it to X ($e(X)$ increases, $e(Y) \geq |Y| - 2$ and $e(X, Y)$ does not increase). So the minimum degree in Y is at least three (but the underlying minimum degree may be one). This implies that $e(Y) \geq 3|Y|/2$. Let $Y := \{y_1, \dots, y_{|Y|}\}$ where the

vertices are indexed in the increasing order according to their degree in Y . For every $i \neq |Y|$, we have $e(Y) \geq (3(|Y| - 2) + d_Y(y_i) + d_Y(y_{|Y|}))/2$. Furthermore,

$$\begin{aligned} e(Y \setminus y_i) &\geq (3(|Y| - 2) + d_Y(y_i) + d_Y(y_{|Y|}))/2 - d_Y(y_i) \\ &\geq 3(|Y| - 2)/2 \\ &\geq |Y| - 2 \\ &\geq b - 2. \end{aligned}$$

We now discuss the two different cases.

- Assume that $e(Y \setminus y) \geq b - 2$ for every $y \in Y$. We denote by Y' the (nonempty) set of vertices of Y with at least one neighbour in X . Set $Y'' := Y \setminus Y'$. Denote by c the minimum excess of a vertex of Y' . Observe that every vertex of Y'' has degree at least four in Y . Thus summing the degrees of the vertices of Y gives

$$2e(Y) \geq e(X, Y) + 4|Y''| + c|Y'| \quad (1)$$

Let $y \in Y'$ such that $\text{exc}(y) = c$. Since the partition $(X \cup y, Y \setminus y)$ is not an $(a + 1)$ -partition, we have $e(X, Y) + c > (2m - n)/3 + 1$. Since $m = e(X, Y) + e(X) + e(Y)$, this implies

$$e(X, Y) + 3c > 2e(X) + 2e(Y) - n + 3 \quad (2)$$

Equations (1) and (2) give:

$$e(X, Y) + 3c > 2e(X) + e(X, Y) + 4|Y''| + c|Y'| - n + 3 \quad (3)$$

Since $e(X) \geq n - |Y| - 1$, we get $3c > n - 2|Y| + 4|Y''| + c|Y'| + 1$. So $3c > n + 2|Y| + (c - 4)|Y'| + 1$, and finally $n + 2|Y| < (c - 4)(3 - |Y'|) + 11$. If $c = 4$, we get $n + 2|Y| \leq 10$, impossible. If $c = 3$, we get $n + 2|Y| - |Y'| \leq 7$, impossible. If $c = 2$, we get $n + 2|Y| - 2|Y'| \leq 4$, impossible. If $c = 1$, we get $n + 2|Y| - 3|Y'| \leq 1$, which can only hold if $|Y| = |Y'| = n - 1$. Thus, X consists of a single vertex, completely joined to Y , against the fact that G is triangle-free. Finally $c > 4$, and consequently $|Y'| < 3$. Observe that $|Y'| > 1$ since G is 2-connected. Thus $|Y'| = 2$. Let y_1, y_2 be the vertices of Y' , indexed in such a way that $e(y_1, X) + e(y_2, Y'') \geq e(y_2, X) + e(y_1, Y'')$. Let $X_1 := X \cup y_1$ and $Y_1 := Y \setminus y_1$. Since $y_1 \in Y'$, we have that $e(X_1) \geq a$. Moreover $e(Y_1) \geq b - 2$. Observe also that $e(y_1, y_2) \leq e(Y \setminus \{y_1, y_2\})$: this is obvious if $e(y_1, y_2) = 0$, and if there is an edge between y_1 and y_2 , since G is triangle-free with minimum degree four, the minimum degree in $Y \setminus \{y_1, y_2\}$ is at least two. So

$$e(Y \setminus \{y_1, y_2\}) \geq |Y| - 2 \geq \lfloor n/2 \rfloor - 1 \geq e(y_1, y_2).$$

In all $e(X_1, Y_1) \leq e(X_1) + e(Y_1)$. And since $n - 2 \leq e(X_1) + e(Y_1)$, we have $e(X_1, Y_1) \leq 2e(X_1) + 2e(Y_1) - n + 2$, which implies $e(X_1, Y_1) \leq (3m - n)/2 + 1$. So the partition (X_1, Y_1) is good.

- Now assume that there exists a vertex $y \in Y$ such that $e(Y \setminus y) \leq b - 3$. We denote by Y' the set of vertices of $Y \setminus y$ with at least one neighbour in X . Set $Y'' := Y \setminus (Y' \cup y)$. Observe that since every vertex of Y'' has underlying degree four in Y , we have $e(Y \setminus y) \geq 3|Y''|/2$. Thus, $|Y''| \leq (2|Y| - 6)/3$. Since $|Y| \geq 3$, we have $|Y''| \leq |Y| - 3$, and finally $|Y'| \geq 3$. Denote by c the minimum excess of a vertex of Y' . Summing the degrees of the vertices of Y gives $2e(Y) \geq e(X, Y) + 4|Y''| + c|Y'| + exc(y)$. Equation (2) still holds, so

$$exc(y) < 3c + n - 3 - 2e(X) - 4|Y''| - c|Y'| \leq 3c - 1 - e(X) - 3|Y''| - (c - 1)|Y'|$$

since $e(X) + |Y''| + |Y'| \geq n - 2$. Therefore $exc(y) < -e(X) - 3|Y''| - (c - 1)(|Y'| - 3) + 2$. Since $|Y'| \geq 3$ and $c \geq 1$, we have $exc(y) \leq 1 - e(X)$. Moreover, since the minimum degree in Y is at least three, summing the degrees in Y of the vertices of $Y \setminus y$ gives that $3(|Y| - 1) \leq 2e(Y \setminus y) + d_Y(y) \leq 2b - 6 + d_Y(y)$. Finally, $d_Y(y) \geq |Y| + 3$ and by the fact that $exc(y) \leq 1 - e(X)$, we have $d_X(y) \geq |Y| + e(X) + 2$. Recall that X contains a vertex x with maximum degree in G . In particular both x and y have degree at least $2|Y| + e(X) + 5$. Observe that $d_Y(x)$ is at least $2|Y| + 5$. Now the end of the proof is straightforward, it suffices to switch x and y to obtain the good partition $(X_1, Y_1) := ((X \cup y) \setminus x, (Y \cup x) \setminus y)$. The only fact to care of is $e(x, y)$. Indeed if $e(x, y)$ is at most $e(X)$, we have:

1. $e(Y_1) \geq d_{Y_1}(x) \geq 2|Y| + 5 - e(x, y) \geq 2|Y| - e(X) \geq |Y| \geq n/2$.
2. $e(X_1) \geq d_{X_1}(y) \geq |Y| + e(X) + 2 - e(x, y) \geq n/2$.
3. Finally, since the excess of y is at most $1 - e(X)$, we have $d_X(y) \geq d_Y(y) + e(X) - 1$, hence $d_{X_1}(y) \geq d_Y(y) - 1$. Moreover $d_{Y_1}(x) \geq 2|Y| + 5 - e(X) \geq e(X) + 5 \geq d_X(x) + 5$. In all, we have $e(X_1, Y_1) \leq e(X, Y) - 4 \leq (2m - n)/3 + 1$, since (X, Y) is an a -partition.

To conclude, we just have to show that $e(x, y)$ is at most $e(X)$. Assume for contradiction that $e(x, y) \geq a$. We consider the partition into $X_2 := \{x, y\}$ and $Y_2 := V \setminus \{x, y\}$. Observe that the minimum underlying degree in Y_2 is at least two, since G being triangle-free, a vertex of Y_2 can only be joined to at most one vertex of X_2 . Thus $e(Y_2) \geq b - 2$. By maximality of a , (X_2, Y_2) is not an $(a + 1)$ -partition, thus $e(X_2, Y_2) > (2m - n)/3 + 1$, that is $d(x) + d(y) - 2e(x, y) > (2m - n)/3 + 1$.

Consider now any partition (X_3, Y_3) such that $(x \cup N(x)) \setminus y \subseteq X_3$ and $(y \cup N(y)) \setminus x \subseteq Y_3$. We have $e(X_3) \geq d(x) - e(x, y) \geq n/2$. Similarly $e(Y_3) \geq n/2$. So, if this partition is not good, we must have $e(X_3, Y_3) > (2m - n)/3 + 1$. Thus $m - (d(x) + d(y) - 2e(x, y)) > (2m - n)/3 + 1$. This gives $m > 2(2m - n)/3 + 2$, and finally $m < 2n - 6$ which is impossible since the minimum degree in G is at least four. ■

5 Conclusion

Note that the only case where $\text{bw}(G) > \text{bw}(M_G)$ is when G has a bridge e and that $\text{bw}(M_G) = 1$, that is when G is a tree that is not a star.

A consequence of our result is a new proof of the fact that the branch-width of a connected planar graph that is not a tree and the branch-width of its dual are the same, for previous proofs see [4] and [1]. Indeed, if r_M is the rank function of a matroid, the rank function r_{M^*} of the dual matroid is such that $r_{M^*}(U) = |U| + r_M(E \setminus U) - r_M(E)$ which implies that $w_M(E_1, E_2) = w_{M^*}(E_1, E_2)$ and that $\text{bw}(M) = \text{bw}(M^*)$. The result follows from the fact that if G is a planar graph and M_G its graphic matroid, the dual matroid of M_G is M_{G^*} .

Note that this dual property also holds for stars and thus the only planar graphs G such that $\text{bw}(G) = \text{bw}(G^*)$ are exactly the planar graphs such that $\text{bw}(G) = \text{bw}(M_G)$. We feel that the natural definition for the branch-width of graphs is the matroidal one.

An independent proof of the equality of branchwidth of cycle matroids and graphs was also given by Hicks and McMurray [3]. Their method is based on matroid tangles and is slightly more involved than ours.

References

- [1] V. Bouchitté, F. Mazoit and I. Todinca, Chordal embeddings of planar graphs, EuroComb'01 (Barcelona), *Discrete Math.*, **273** (2003), 85–102.
- [2] J.F. Geelen, A.M.H. Gerards, N. Robertson and G.P. Whittle, On the excluded minors for the matroids of branch-width k , *J. Combin. Theory Ser. B*, **88** (2003), 261–265.
- [3] I.V. Hicks and N.B. McMurray, The branchwidth of graphs and their cycle matroids, preprint.
- [4] D. Lapoire, Treewidth and duality for planar hypergraphs, preprint.
- [5] N. Robertson and P. Seymour, Graph minors. X. Obstructions to tree-decomposition, *J. Combin. Theory Ser. B*, **52** (1991), 153–190.